

HOW TO PUT A HEAVIER HIGGS ON THE LATTICE[§]

by

Urs M. Heller[†], Herbert Neuberger[‡], and Pavlos Vranas[†][†] *Supercomputer Computations Research Institute
The Florida State University
Tallahassee, FL 32306*[‡] *Department of Physics and Astronomy
Rutgers University
Piscataway, NJ 08855-0849***Abstract**

Lattice work, exploring the Higgs mass triviality bound, seems to indicate that a strongly interacting scalar sector in the minimal standard model cannot exist while low energy QCD phenomenology seems to indicate that it could. We attack this puzzle using the $1/N$ expansion and discover a simple criterion for selecting a lattice action that is more likely to produce a heavy Higgs particle. Our large N calculation suggests that the Higgs mass bound might be around 850 GeV , which is about 30% higher than previously obtained.

[§] Submitted to Physical Review Letters

The recent quantitative results that emerged from the investigation of triviality in the $\lambda(\vec{\phi}^2)^2$ theory on the lattice [1] have given an upper bound on the ratio of the Higgs mass, m_H , to the weak scale, $f_\pi = 246 \text{ GeV}$, of about 2.5–2.7. In the hypothetical world of QCD with just the up-down doublet of quarks with zero masses, which is described by the same effective theory at low energies, one believes, on phenomenological grounds, that the above ratio would be significantly larger, equal to about 6–8. This is admittedly an extreme situation because the sigma enhancement in $\pi\pi$ scattering doesn't quite qualify to be called a resonance, and, it comes accompanied by a host of other resonances, representing all kinds of nontrivial “cutoff effects”. Still, there is a large gap between 2.7 and 6 and we are left with essentially two alternatives: either the QCD example provides a totally unreasonable estimation for the bound, or, we admit that the lattice result, while solidly established within a subclass of actions, is relatively far from a realistic estimate for the bound. We would have much less cause for worry if the lattice bound for the ratio came out to be about 4 (this would correspond to $m_H \simeq 1 \text{ TeV}$) or more [2].

This issue could be addressed in principle by an extensive search of all possible lattice actions. In practice one needs some theoretical indication as to what kind of regularized action is likely to produce a higher bound. If the physical picture behind the particular form of the suggested action is reasonable, an investigation of its consequences would help resolve the above puzzle by either refuting the existing lattice bound or by strengthening the contention that the QCD example is grossly exaggerated.

The purpose of the present note is to make a proposal for such a generalized action and sketch the theoretical basis for it. The basic point of the proposal is that one should put on the lattice nonlinear $O(4)$ models that have a naive continuum limit with freely adjustable four derivative couplings and study them nonperturbatively. We solve the model in the large N limit of $O(N)$ and find the region in coupling space where the highest bound is expected; we test the extent to which the results are dependent on the regularization and extract the basic features that are generic. The results can be explained by a rough physical picture as follows: When the regularized model is nonlinear one has to think about the Higgs resonance as a loosely bound state of two pions in an $I = 0, J = 0$ state. Pions in such a state attract because superposing the field configurations corresponding to individual pions makes the state look more like the vacuum and hence lowers the energy. The four derivative terms in the action can add to this attraction; if the attraction is increased to a certain level the bound state will become massless and stable. This occurs while the system is still relatively deep in the broken phase; hence the weak scale f_π is nonvanishing. The appearance of this massless bound state signals a new kind of phase transition; the ordinary phase transition occurs when both f_π and m_H vanish and leads to a scale invariant theory at the transition - here we still have a scale and therefore scale invariance seems to be spontaneously broken, with the massless boundstate, the Higgs particle, playing the role of the expected dilaton. This phenomenon is reminiscent of the results obtained in three dimensions by Bardeen, Moshe and Bander [3]. As far as the Higgs mass bound goes, it is clear that we have to try to be as far away as possible from this new transition point; we wish to induce enough repulsion between the pions to delay the creation of the bound state as much as possible. Of course, the interaction is still attractive overall, because only the order p^4 term in the interaction can be tampered with; the order p^2 and $p^4 \log(p^2)$ terms are fixed by current algebra and are attractive [4]. Moreover, in the critical regime the model still is governed by the $\lambda(\vec{\phi}^2)^2$ trivial fixed point and is expected therefore to generate a one particle state corresponding to the σ field component. While it is quite possible that the above described scale invariance breaking transition cannot actually materialize at $N = 4$ (we shall see that at $N = \infty$ it does appear in all regularizations that we tested), the basic trend for the magnitude of the triviality bound must be the same.

Assuming global $O(4)$ symmetry and that the leading cutoff effects are represented by dimension six operators, one arrives, after eliminating couplings that won't lead to observable effects in the S-matrix to leading order in the inverse cutoff, to a space of actions that depends on 4 parameters [5]. One may try to induce the dependence on the associated renormalized operators by simply writing identical forms for terms in the bare action. However, we are looking for a bound on the physical four point coupling and it is likely that it obtains in the limit that the bare four point coupling is taken to infinity. When the bare coupling becomes infinite the bare model is reduced to a nonlinear model; this has the effect that the naive forms of the dimension six operators in terms of bare linear fields become trivial. The parameterization in terms of linear fields and operators of naive dimension six breaks down completely at the point where the bare theory becomes nonlinear (we have tested this explicitly in the linear model, within $1/N$, and with a particular form

of a bare dimension-six term in the Lagrangian). So we were expecting to be left with three freely adjustable parameters (within some range) but we ended up with only one. This is the case that has been investigated nonperturbatively until now [1]. To include the effects of dimension six operators at physical scales we need to put in operators of higher order in derivatives in the bare, nonlinear, theory that describes the physics at cutoff scales. There are exactly two additional terms that one can write down if one restricts the number of derivatives to four [4]; thus the right number of parameters is obtained. This time, unlike in the linear case, the effect of the bare couplings has no reason to disappear.

It is well known that the $1/N$ expansion, at least to leading order, doesn't have to work well quantitatively simply because there is a factor of $N + 8$ in the coefficients of the β function and therefore $N = 4$ is badly approximated by $N = \infty$ [6-8]. We shall say more about this later; for the moment keep in mind that we are interested mainly in relative effects, that is, in how the bound varies with the action, and less in the absolute magnitude of the bound. To get the latter one would have to resort to numerical means.

If a strongly interacting scalar sector could be produced on the lattice its investigation would require some nontrivial extension of the tools that were used until now. However, if we only wish to establish the existence of a strongly interacting regime, we can do that indirectly, while restricting our attention just to weak couplings. For example, a strongly coupled Higgs sector was excluded on lattices with a pure nearest-neighbor coupling, by showing that, for a ratio m_H/f_π of only 2.5, one already had m_H/Λ equal to 0.5, leaving no "room" below the cutoff Λ for further growth in the coupling; if we could show that a different action would, at the same value of m_H/f_π , give m_H/Λ equal to 1/10 say, our case would be made. Therefore, our subsequent study is restricted to the weakly coupled sector. Generically, we expect there that

$$\frac{m_H}{\Lambda} \approx C(b) \exp[-16\pi^2 f_\pi^2 / (Nm_H^2)], \quad (1)$$

where $C(b)$ is a function of the additional parameters we introduced in the action. We aim to show that by varying b $C(b)$ can be significantly lowered below the value it had when no four derivative terms were included (this is the only case that has been investigated to date). Our finding is best summarized in a qualitative self-explanatory diagram that we checked for a class of Pauli-Villars regularizations and for the F_4 lattice regularization (the generalization of the calculations to the hypercubic lattice should be a simple exercise).

We first look at the models with Pauli-Villars, or higher derivative, regularization. We choose to replace the ordinary propagator by $1/[p^2(1 + (p^2/M^2)^n)]$ where M is a "bare" cutoff (as opposed to Λ the physical cutoff to be defined below) and $n \geq 3$ is an integer. The limit $n = \infty$ corresponds to a sharp momentum cutoff and is expected to be singular. For any finite n the bare propagator has n "ghost" poles all situated on the unit circle in the complex s/M^2 plane. We shall define the physical cutoff, Λ , as the square root of the modulus of the location of the closest complex pole in s in the full pion propagator (at leading order in $1/N$). The action, written in Euclidean space, is:

$$S = \int_x \left[\frac{1}{2} \vec{\phi} g(-\partial^2) \vec{\phi} - \frac{b_1}{2N} (\partial_\mu \vec{\phi} \cdot \partial_\mu \vec{\phi})^2 - \frac{b_2}{2N} (\partial_\mu \vec{\phi} \cdot \partial_\nu \vec{\phi} - \frac{1}{4} \delta_{\mu,\nu} \partial_\sigma \vec{\phi} \cdot \partial_\sigma \vec{\phi})^2 \right] \quad (2)$$

where $g(-\partial^2) = (-\partial^2)[1 + (-\partial^2/M^2)^n]$, and the field is subjected to the constraint $\vec{\phi}^2 = N\beta$. The partition function is $\mathcal{Z} = \int [d\vec{\phi}] \exp[-S]$.

Using the technique employed in ref. [9] we find the phase diagram of the model paying particular attention to the competition between various saddles. We set

$$\vec{\phi} = \sqrt{N} \vec{v} + \hat{v} H(x) + \vec{\pi}(x) \quad (3)$$

with $\hat{v} = \vec{v}/v$, $v = |\vec{v}|$, $\hat{v} \cdot \vec{\pi} = 0$, $\int_x H(x) = 0$, $\int_x \vec{\pi}(x) = 0$. We define the pion wave function renormalization constant Z_π for the field $\vec{\pi}$ as usual. The pion decay constant f_π can be easily gotten from v and Z_π .

It turns out that it is more convenient, for the broken phase, to parameterize the theory by Z_π :

$$b_1 = \left(\frac{4\pi}{M^2} \right)^2 \frac{n \sin(2\pi/n)}{2\pi} \frac{Z_\pi - 1}{Z_\pi^{2(n-1)/n}} \quad (4)$$




Fig. 1 : **Summary Diagram**

Qualitative Features Common to all Models Investigated

where $0 \leq Z_\pi \leq 2(n-1)/(n-2)$. In this range, b_1 increases monotonically with Z_π and at the larger end-point we have $\partial b_1 / \partial Z_\pi = 0$. The portion of the phase diagram we are interested in does not depend on b_2 because giving an expectation value to the $O(N)$ singlets associated with b_2 would break Lorentz invariance. The other coupling is best replaced by the unitless weak scale, $F = f_\pi / \Lambda$. The physical cutoff, Λ , is given by $\Lambda = M / Z_\pi^{1/(2n)}$. From the large N saddle point equations we obtain the expectation value v which is related to f_π by $v^2 = Z_\pi f_\pi^2$ resulting in:

$$\frac{(4\pi F)^2}{N} = \left(\frac{4\pi}{M}\right)^2 \beta Z_\pi^{(1-n)/n} - \frac{\pi}{n \sin(\pi/n)}. \quad (5)$$

There is a second order transition line at $F = 0$ and, on it, at the end of the allowed range of Z_π , there is a tricritical point where the second order line turns into first order. At the tricritical point one obtains a massless pole in the two point correlation function of the trace of the energy-momentum tensor. We shall not be concerned with the region beyond the tricritical point. The calculation of m_H proceeds in the standard manner; it turns out that one has to find the complex zeros of the determinant of a four by four matrix. Since we restrict our attention to the perturbative regime, i.e. $F, p \ll 1$ we can replace the quantity m_H by the technically more accessible real quantity m_R , defined from the smallest positive zero in $s = -p^2$ of the real part of the determinant by $m_R = \sqrt{s}$. We are really interested in $M_R = m_R / \Lambda$. If we also take $M_R \ll 1$ of the same order (up to logs) as F , the fourth row and column of the matrix decouple and the

dependence on b_2 disappears. We need the expression for the “bubble” diagram

$$\begin{aligned} B_{PV}(p) &= \int \frac{d^4 k}{(2\pi)^4} \frac{1}{g(k-p/2)g(k+p/2)} \\ &= -\frac{1}{16\pi^2} \left[\log(p^2/M^2) - \frac{n-1}{n} \right] + O(p^2/M^2) \end{aligned} \quad (6)$$

which holds for any $n < \infty$ (for $n = \infty$ the correction term goes as $\sqrt{p^2/M^2}$ and represents a non-local term in the effective action as expected when employing a sharp momentum cutoff). One has then:

$$\begin{aligned} M_R^2 &\approx \exp \left[-\frac{2}{N} \left(\frac{4\pi F}{M_R} \right)^2 \right] C^2(Z_\pi), \\ C(Z_\pi) &= \exp \left[\frac{n-1}{2n} \left(1 - \frac{\pi}{n \tan(\pi/n)} \frac{1-Z_\pi}{1-Z_\pi(n-2)/(2(n-1))} \right) \right]. \end{aligned} \quad (7)$$

There is no four derivative term when $Z_\pi = 1$; the minimal C obtains at $Z_\pi = 0$ which is as far away from the tricritical point as possible:

$$\frac{C(Z_\pi = 1)}{C(Z_\pi = 0)} = \exp \left[\left(\frac{n-1}{2n} \right) \frac{\pi}{n \tan(\pi/n)} \right] \leq \exp(0.5) \approx 1.65. \quad (8)$$

We have obtained some increase in the cutoff but not a factor of about 5 as we would have liked. However, we know from the work of Einhorn [7] that even at $b_1 = 0$ this particular large N model will have a strongly interacting sector; therefore it may be that there is no room left to dramatically increase the cutoff even further. It is also clear that if it turned out that the lattice actions that were so far investigated really correspond to an action with $b_1 > 0$ in Pauli-Villars regularization, a much larger variation could be obtained because C diverges at some finite, positive b_1 . $b_1 > 0$ means that the extra term has induced more attraction in the $I, J = 0$ channel and the divergence in C occurs at the tricritical point where the resonant bound state becomes massless and stable.

We turn now to the lattice. Since our objective here is to provide a useful indication for what kind of actions would be instructive to study by nonperturbative means, we pick the lattice regularization that is best suited for this purpose, namely we put the theory on an F_4 lattice. This lattice has a larger symmetry group than the more commonly used hypercubic lattice which ensures that no Lorentz invariance breaking four derivative terms appear in the (naive) continuum limit of its lattice actions. The F_4 lattice can be viewed as a hypercubic lattice (Z^4) with unit lattice spacing from which all the sites with an odd sum of coordinates have been erased [5]. We shall denote sites on F_4 by x, x', x'' , and links by $\langle x, x' \rangle, l, l'$. The action is chosen as

$$\begin{aligned} S &= -2N\beta_0 \sum_{\langle x, x' \rangle} \vec{\Phi}(x) \cdot \vec{\Phi}(x') - N\beta_1 \sum_{\langle x, x' \rangle} \left[\vec{\Phi}(x) \cdot \vec{\Phi}(x') \right]^2 \\ &\quad - N\frac{\beta_2}{48} \sum_x \left[\sum_{\substack{l \cap x \neq \emptyset \\ l = \langle x, x' \rangle}} \vec{\Phi}(x) \cdot \vec{\Phi}(x') \right]^2. \end{aligned} \quad (9)$$

Here the field is constrained by $\vec{\Phi}^2(x) = 1$. To express the action in a form closer to the one we employed before we rescale the field $\vec{\phi} = \sqrt{6N(\beta_0 + \beta_1 + \beta_2)} \vec{\Phi}$ (we only consider the region $\beta_0 + \beta_1 + \beta_2 > 0$). The action has the same form as above (eq. (2)), up to terms that contain fourth order derivatives, if one identifies:

$$\begin{aligned} g(k_\mu) &= \frac{1}{6} \sum_{\mu \neq \nu} [2 - \cos(k_\mu + k_\nu) - \cos(k_\mu - k_\nu)] \approx k^2 - \frac{(k^2)^2}{12} + O(k_\mu^6), \\ b_1 &= \frac{1}{48} \frac{\beta_1 + \beta_2}{(\beta_0 + \beta_1 + \beta_2)^2}, \quad b_2 = \frac{1}{36} \frac{\beta_1}{(\beta_0 + \beta_1 + \beta_2)^2}. \end{aligned} \quad (10)$$

Again b_2 plays no role and one may set β_1 to 0. The situation is essentially the same as in the Pauli-Villars case. b_1 can be traded for the pion wave function renormalization constant,

$$b_1 = \frac{1}{Z_\pi} - \frac{1}{Z_\pi^2}, \quad (11)$$

where Z_π varies between 0 and 2. The standard non-linear action corresponds again to $Z_\pi = 1$ and, on the line where the pion decay constant F vanishes, $Z_\pi = 2$ corresponds to the tricritical point. F is given by:

$$\frac{F^2}{N} = \frac{6(\beta_0 + \beta_1 + \beta_2)}{Z_\pi} - r_0, \quad r_0 = \int_k \frac{1}{g(k)} = 0.13823. \quad (12)$$

The momentum integral is over the F_4 Brillouin zone and a factor of $1/(2\pi)^4$ has been absorbed in the definition of $\int_k$; the numerical value is taken from ref. [5].

We proceed now to find M_R defined exactly as before; for this we need the lattice “bubble” which can be easily evaluated using the results of ref. [5] (but note a missing minus sign in the value of r_1 in equation (4.12) there):

$$B_{F_4}(p) = -\frac{1}{16\pi^2} \log(p^2) + c_1 + c_2 p^2 + O(p^4 \log p^2), \quad (13)$$

$$c_1 = 0.0466316, \quad c_2 = 0.0005497.$$

Unlike in the Pauli-Villars case, the cutoff (defined for example by a certain amount of violation of Euclidean rotational invariance in the full pion propagator) is independent of the couplings in the action.

This time we shall be interested to go beyond the $O(1)$ term and that’s why we evaluated the next correction in p to the bubble (13). Ignoring this correction for the moment we get the usual formula for M_R (eq. (7)), but the coefficient C is now given by,

$$C(Z_\pi) = \exp \left[8\pi^2 \left(c_1 - r_0^2 \frac{1 - Z_\pi}{1 - Z_\pi/2} \right) \right], \quad (14)$$

leading to:

$$\frac{C(Z_\pi = 1)}{C(Z_\pi = 0)} = \exp(8\pi^2 r_0^2) \approx 4.521. \quad (15)$$

We see that we have obtained a much larger variation.

In practice, for vectorization purposes, it would be better to have an action that involves only nearest neighbors on the F_4 lattice. Nothing in what we have said until now changes if we replace the β_2 term in the action by

$$N \frac{3\beta_2}{8} \sum_x \sum_{\substack{\langle uv \rangle \\ l, l' \cap x \neq \emptyset, \ l \cap x' \neq \emptyset, \ l' \cap x'' \neq \emptyset, \ x, x', x'' \text{ all n.n.}}} \left[\left(\vec{\Phi}(x) \cdot \vec{\Phi}(x') \right) \left(\vec{\Phi}(x) \cdot \vec{\Phi}(x'') \right) \right]. \quad (16)$$

To check sensitivity to higher order terms in F^2 and/or M_R^2 , while still neglecting b_2 , we go to the next higher order in F, p, M_R when we find the Higgs mass. It turns out that the numerical correction is small (of the order of 1–2 %) even for $M_R \approx 1$. For the usually studied action with b_1 and $b_2 = 0$ we can check how well the $1/N$ expansion works, on the quantitative level, in the region of main interest, namely $m_R/f_\pi \approx 2.0 - 2.5$; in this region we have numerical data at $N = 4$ (see last reference in [1]). We find agreement to within 25 percent for the M_R/F ratio as a function of M_R ; the large N results are systematically larger than the $N = 4$ ones but the relative change in the ratio M_R/F when M_R is varied is more accurately predicted. This is enough to convince us that our large N results ought to be taken seriously as an indication for what is going to happen in the real system. For example, we would predict that if we limit cutoff effects by the reasonable [5] requirement $M_R \leq \sqrt{2}/2$ we find, with $N = 4$:

$$\frac{M_R}{F} \leq \frac{2\pi}{\sqrt{\log [\sqrt{2}C(Z_\pi)]}}. \quad (17)$$

This gives for the nearest neighbor case ($Z_\pi = 1$) $M_R/F = 3.1$ while for the extreme case ($Z_\pi = 0$) one finds $M_R/F = 4.0$. For $N = 4$ the numerical result is smaller by about 20%. We therefore are led to conjecture that for the simplest case with $\beta_1 = 0$ the bound would increase by 30% to about 850 GeV.

We should comment here on the possible relationship between this note and the work based on effective Lagrangians that was carried out in the context of the investigation of the scalar sector of the standard model by several groups recently [10]. Although in both cases one employs a nonlinear Lagrangian of identical forms (at least for $N = 4$) there is a fundamental difference that ought to be stressed: In the effective Lagrangian approach one uses *renormalized* perturbation theory as defined by Weinberg [4] to summarize all relations between soft pion amplitudes to any finite order in the external momenta of the pions; one then tries to extrapolate [10] these results by several schemes, all attempting to approximately impose unitarity and crossing, to the region of energies dominated by the tails of the lowest resonances. At no point does this approach consider the question of whether the physics so defined can be consistently extended to the cutoff (see second reference in [4] for some discussion). Technically speaking, chiral renormalized perturbation theory, very much like ordinary perturbation theory, does not provide the relation between the parameters in the bare Lagrangian (i.e. cutoff scale physics) and the parameters that describe low energy properties; it only provides a very large number of constraints between all the low energy physical processes. Thus, questions having to do with triviality are completely outside the reach of these approaches. In more intuitive terms, we are taking here very seriously a model defined by a chiral Lagrangian, all the way to the highest energies, and seek for constraints on the allowable low energy effects that result from the requirement that they occur below the cutoff. If the cutoff is very high relative to the lowest resonance we are bound, by the general principles of the renormalization group, to view this as just another regularization of $\lambda(\phi^2)^2$ and hence a perfectly reasonable framework for investigating consequences of triviality.

We have not discussed in any detail the term with coupling b_2 . We have seen that it has no effect in the perturbative regime; however, when terms beyond the equivalent of the $O(p^2)$ correction to the bubble are taken into account, b_2 comes into play and can become important. The structure of the term indicates that b_2 is the right dial to tune if we wish to make our theory look more like QCD. This suggests that it might help to make the Higgs mass bound even higher. But in the present note we were interested mainly in estimating the Higgs-mass triviality bound when nothing special (technicolor-like) has been imposed on the theory. Further discussion of the b_2 term would take us too far afield. We shall also omit any discussion of subleading terms in the $1/N$ expansion; particularly interesting would be the corrections to Z_π and to the phase structure in the vicinity of the tricritical point. These topics and a comprehensive analysis of Renormalization Group flows in the models under consideration will hopefully be addressed in future work.

We should also comment here on the fact that it is well known that once the scalar sector is strongly interacting, looking at m_R rather than m_H , is a bad approximation due to the large width [11]. This effect leads to a saturation of the growth of m_H much before m_R stops growing and can be best seen in the $1/N$ expansion, as shown by Einhorn [7]. It is important to realize that this sort of bound has nothing to do with triviality; it appears even when the issue of triviality is totally ignored, for example, when the bare four point coupling in the linear model is allowed to become negative and the theory has a tachyonic instability. Our calculations can all be done for a relatively narrow Higgs, and all they aim to show is that a strongly interacting scalar sector in the minimal standard model has not yet been convincingly ruled out. The relation between m_R and m_H alluded to above may very well hold in such a sector, if only we could show that one exists.

In summary, we propose that the scalar sector of the minimal Standard Model can be made to have a stronger renormalized self coupling by introducing, at the cutoff level, four-derivative couplings that induce some repulsion between Goldstone Bosons. Within the $1/N$ expansion this increases the Higgs mass bound, compared to the usual lattice result, by up to about 30%. Obviously it would be extremely interesting to study, by numerical means, whether this conclusion persists for the physically relevant case of $N = 4$. This would help resolve the puzzle presented at the beginning of this Letter. We hope to address this question in the future.

ACKNOWLEDGEMENTS.

This research was supported in part by the Florida State University Supercomputer Computations Research Institute which is partially funded by the U.S. Department of Energy through DOE grant # DE-FG05-85ER250000 (UMH and PV) and under DOE grant # DE-FG05-90ER40559 (HN). UMH also

acknowledges partial support by the NSF under grant # INT-8922411 and he would like to thank F. Karsch and the other members of the Fakultät für Physik at the University of Bielefeld for the kind hospitality while part of this work was done. HN would like to thank F. David and M. Einhorn for useful discussions.

REFERENCES

- [1] M. Lüscher and P. Weisz, *Phys. Lett.*, **B212** (1988) 472; J. Kuti, L. Lin and Y. Shen, *Phys. Rev. Lett.*, **61** (1988) 678; A. Hasenfratz, K. Jansen, J. Jersák, C. B. Lang, T. Neuhaus, H. Yoneyama, *Nucl. Phys.*, **B317** (1989) 81; G. Bhanot, K. Bitar, *Phys. Rev. Lett.*, **61** (1988) 427; G. Bhanot, K. Bitar, U. M. Heller, H. Neuberger, *Nucl. Phys.*, **B353** (1991) 551.
- [2] see the *The Standard Model Higgs Boson* edited by M. B. Einhorn, *CPSC* Vol. **8**, Elsevier Pub.
- [3] W. Bardeen, M. Moshe, M. Bander, *Phys. Rev. Lett.*, **52** (1983) 1188.
- [4] S. Weinberg, *Physica*, **96A** (1979) 327; H. Georgi, *Weak Interactions and Modern Particle Theory* (1984) Addison-Wesley Pub.
- [5] G. Bhanot, K. Bitar, U. M. Heller, H. Neuberger, *Nucl. Phys.*, **B343**, (1990) 467.
- [6] R. Dashen, H. Neuberger, *Phys. Rev. Lett.*, **50** (1983) 1897.
- [7] M. Einhorn, *Nucl. Phys.*, **B246** (1984) 75.
- [8] L. Lin, J. Kuti, Y. Shen, in *Lattice Higgs Workshop, FSU, May 16–18 1988*, Editors B. Berg., et. al.
- [9] F. David, D. Kessler, H. Neuberger, *Nucl. Phys.*, **B257** [FS14] (1985) 695.
- [10] J. Bagger, S. Dawson, G. Valencia, BNL – 45782, Brookhaven Nat. Lab. preprint; S. Dawson, G. Valencia, *Nucl. Phys.*, **B352** (1991) 27; J. F. Donoghue, C. Ramirez, G. Valencia, *Phys. Rev.*, **D39** (1989) 1947; A. Dobado, M. Herrero, J. Terron, CERN–TH.5670/90, CERN preprint and references therein.
- [11] B. W. Lee, C. Quigg, H. B. Thacker, *Phys. Rev.*, **D16** (1977) 1519.